


# Substance Abuse Trends in Maine

Epidemiological Profile 2015

## Downeast District



Department of Health  
and Human Services

Maine People Living  
Safe, Healthy and Productive Lives

Paul R. LePage, Governor

Mary C. Mayhew, Commissioner

Produced for Maine Department of Health and Human Services  
Office of Substance Abuse and Mental Health Services  
by Hornby Zeller Associates, Inc.  
October 2015



# **Substance Abuse Trends in Maine**

**Epidemiological Profile 2015**

## **Downeast District**

**THIS REPORT IS PRODUCED FOR  
THE MAINE OFFICE OF SUBSTANCE ABUSE  
COMMUNITY EPIDEMIOLOGY SURVEILLANCE NETWORK  
WITH SUPPORT FROM THE PARTNERSHIPS FOR SUCCESS GRANT THROUGH THE U.S.  
DEPARTMENT OF HEALTH AND HUMAN SERVICES  
SUBSTANCE ABUSE AND MENTAL HEALTH ADMINISTRATION**

**Produced by  
Hornby Zeller Associates, Inc.  
373 Broadway  
South Portland, ME 04106  
October 2015  
(207) 773-9529  
[www.hornbyzeller.com](http://www.hornbyzeller.com)**



## Table of Contents

---

List of Figures .....	ii
Introduction .....	1
Overview of Downeast Public Health District.....	1
Purpose of this Report.....	1
Consumption of Substances.....	3
Alcohol .....	4
Tobacco.....	10
Prescription Drugs .....	13
Other Illegal Drugs.....	17
Consequences Resulting from Substance Use and Abuse.....	20
Substance Use and Pregnancy .....	21
Referral Services .....	23
Criminal Justice Involvement .....	25
Driving Under the Influence .....	33
Hospital Visits Related to Substance Use .....	37
Overdoses and Related Deaths .....	47
Factors Contributing to Substance Use and Abuse.....	54
Availability and Accessibility .....	55
Perceived Risk and Harm.....	65
Perceived Enforcement.....	71
Mental Health, Suicide and Co-occurring Disorders .....	73
Depression and Anxiety .....	74
Suicide and Suicidal Ideation .....	77
Mental Health and Substance Abuse Co-Occurrence.....	79
Treatment Admissions for Substance Abuse .....	81
Treatment Admissions .....	82
Appendix: Data Sources.....	91
Description of Data Sources .....	91

## List of Figures

---

Figure 1. Percent of high school students by Public Health District who had at least one drink of alcohol during past 30 days: 2009-2013.....	4
Figure 2. Percent of high school students by Public Health District who had at least five drinks in a row during past 30 days: 2009-2013 .....	5
Figure 3. Percent of adults by Public Health District who reported drinking during past 30 days: 2011-2013.....	6
Figure 4. Percent of adults by Public Health District who reported heavy drinking during past 30 days: 2011-2013 .....	7
Figure 5. Percent of adults by Public Health District who reported binge drinking during past 30 days: 2011-2013 .....	8
Figure 6. Percent of adults by Public Health District who reported binge drinking in past 30 days by age group: 2011-13 .....	9
Figure 7. Percent of high school students by Public Health District who reported smoking one or more cigarettes during past 30 days: 2009-2013 .....	10
Figure 8. Percent of high school students by Public Health District who used smokeless tobacco in the past 30 days: 2009-2013 .....	11
Figure 9. Percent of adults by Public Health District who reported smoking a cigarette in the past 30 days: 2011-2013.....	12
Figure 10. Percent of high school students by Public Health District who have taken prescription drugs not prescribed to them by a doctor (past 30 days): 2009-2013.....	13
Figure 11. Misuse of prescription drugs among Maine residents (18 and older) in their lifetime, by gender and public health district: 2011-13 .....	14
Figure 12. Percent of population 18 years old or older who used prescription pain relievers in past year for nonmedical use, by Public Health District: 2010-12 .....	15
Figure 13. Percent of 18 to 25 year olds who used prescription pain relievers in past year for nonmedical use, by Public Health District: 2004-06 to 2010-12 .....	16
Figure 14. Percent of high school students by Public Health District who have used marijuana during past 30 days: 2009-2013 .....	17
Figure 15. Percent of adults who have used marijuana during the past 30 days, by gender and Public Health District: 2012-13.....	18

Figure 16. Percent of high school students by Public Health District that have used cocaine in any form during their lifetime: 2009-2013 .....	19
Figure 17. Number of drug affected baby notifications, by Public Health District: 2010-2013.....	22
Figure 18. Number of 2-1-1 referral calls relating to substance use per 10,000 residents, by Public Health District: 2010-2013 .....	24
Figure 19. Violent crime rate per 10,000, by Public Health District: 2009-2013.....	26
Figure 20. Alcohol-related arrest rate per 10,000, by Public Health District: 2009-2013.....	28
Figure 21. Drug-related arrest rate per 10,000, by Public Health District: 2009-2013 .....	29
Figure 22. Drug-related arrest rate per 10,000, by drug type and Public Health District: 2013.....	30
Figure 23. Maine DEA Drug offense arrests per 10,000 residents, by Public Health District: 2010-2014 .....	31
Figure 24. Maine DEA Drug offense arrests per 10,000 residents, by Public Health District: 2012-14 .....	32
Figure 25. Percent of high school students by Public Health District who reported drinking and driving during the past 30 days: 2009-2013 .....	33
Figure 26. Percent of high school students by Public Health District who rode in a vehicle driven by someone who had taken illegal drugs: 2009-2013 .....	34
Figure 27. Alcohol/Drug-related motor vehicle crash rate per 10,000, by Public Health District: 2010-2014 .....	36
Figure 28. Inpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District: 2010, 2011 .....	38
Figure 29. Inpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District and drug type: 2011 .....	39
Figure 30. Outpatient hospital admissions (per 10,000 people) related to substance use, * by Public Health District: 2010, 2011 .....	41
Figure 31. Outpatient hospital admissions (per 10,000 people) related to substance use*, by Public Health District and drug type: 2011 .....	42
Figure 32. Number of poisonings reported to New England Poison Center 10,000 residents, by intent and Public Health District: 2012-14 .....	44

Figure 33. Number of poisonings reported to New England Poison Center suspected to be substance abuse per 10,000 residents, by Public Health District: 2012-14 .....	45
Figure 34. Number of poisonings reported to New England Poison Center suspected to be suicide attempts per 10,000 residents, by Public Health District: 2012-14 .....	46
Figure 35. Number of overdose EMS responses per 10,000 residents, by Public Health District and primary type of substance involved: 2014 .....	48
Figure 36. Number of overdose EMS responses due to drug/medication per 10,000 residents, by Public Health District: 2011-2014 .....	49
Figure 37. Number of total overdose EMS responses due to alcohol per 10,000 residents, by Public Health District: 2011-2014 .....	50
Figure 38. Naloxone* administrations per 10,000 residents, by Public Health District: 2014.....	51
Figure 39. Drug-related death rate per 100,000, by Public Health District: .....	53
Figure 40. Percent of high school students by Public Health District who reported it was easy to get alcohol: 2009-2013.....	55
Figure 41. Parent perception of accessibility of parent purchased alcohol without parental knowledge: 2013 and 2015 .....	56
Figure 42. Parent perception of teen accessibility of prescription drugs at home without parental knowledge: 2015 .....	57
Figure 43. Percent of high school students by Public Health District who reported it would be easy to get marijuana: 2009-2013.....	58
Figure 44. Number of liquor licensees per 1,000 residents, by Public Health District: 2015.....	59
Figure 45. Dispensed quantity of narcotics per capita, by Public Health District: 2010-2014* .....	61
Figure 46. Dispensed quantity of stimulants per capita, by Public Health District: 2010-2014.....	62
Figure 47. Dispensed quantity of tranquilizers per capita, by Public Health District: 2010-2014 .....	63
Figure 48. Number of poisonings reported to New England Poison Center per 10,000 residents, by drug type and Public Health District: 2013-14 .....	64



Figure 49. Percent of high school students by Public Health District who reported a risk of harm from consuming five or more drinks once or twice per week: 2009-2013.....	65
Figure 50. Percent of population age 12 or older who perceive a great risk from binge drinking, by Public Health District: 2010-12.....	66
Figure 51. Percent of high school students by Public Health District who reported a risk of harm from smoking marijuana regularly: 2009-2013 .....	67
Figure 52. Percent of population age 12 or older who perceive a great risk from smoking marijuana once a month by Public Health District: 2006-2008 to 2010-12* .....	68
Figure 53. Parent disapproval regarding their teen using marijuana: 2013 and 2015 .....	69
Figure 54. Percent of high school students who reported a risk of harm from misusing prescription drugs, by Public Health District: 2009-2013.....	70
Figure 55. Perceived risk among high school students in Downeast PHD of being caught by parents or police for drinking alcohol: 2009-2011 .....	71
Figure 56. Perceived risk among high school students of being caught by police for smoking marijuana, by Public Health District: 2009-2013 .....	72
Figure 57. Percent of adults who have ever been told they have an anxiety disorder, by Public Health District: 2011-12 to 2012-13 .....	74
Figure 58. Percent of adults who have ever been told they have a depression disorder, by Public Health District: 2011-2013 .....	75
Figure 59. Felt sad or hopeless almost every day for two weeks or more in a row during the past year, by Public Health District: 2009-2013.....	76
Figure 60. Number of suicide deaths per 100,000 residents, by Public Health District: 2013-14 .....	77
Figure 61. Percent of high school students who considered suicide during the past year, by Public Health District: 2009-2013 .....	78
Figure 62. Percent of individuals by Public Health District admitted for substance abuse treatment that also had a mental health diagnosis: 2010-2014 .....	79
Figure 63. Number of Maine 2-1-1 referral calls related to mental health services per 10,000 residents, by public health districts: 2010-2014.....	80
Figure 64. Primary drug admissions for in Downeast PHD, by drug type: 2010-2014 .....	82
Figure 65. Primary treatment admissions in Downeast PHD: 2014.....	83

<b>Figure 66. Primary admissions related to alcohol, by Public Health District and drug type: 2010-2014 .....</b>	<b>84</b>
<b>Figure 67. Primary admissions related to synthetic opioids, by Public Health District: 2010-2014 .....</b>	<b>85</b>
<b>Figure 68. Primary admissions related to heroin/morphine, by Public Health District: 2010-2014 .....</b>	<b>86</b>
<b>Figure 69. Primary admissions related to marijuana, by Public Health District and drug type: 2010-2014 .....</b>	<b>87</b>
<b>Figure 70. Secondary treatment admissions in Downeast PHD: 2010-2014 .....</b>	<b>88</b>
<b>Figure 71. Secondary treatment admissions in Downeast PHD, by drug type: 2014.....</b>	<b>89</b>



## Introduction

---

### Overview of Downeast Public Health District

Downeast Public Health District is comprised of two counties, Hancock and Washington, and has a population of 87,035 people. This represents approximately 7 percent of Maine's total population in 2013. Downeast PHD is relatively rural with 21 people per square mile; Hancock has a higher number of persons per square mile than Washington County. Washington is the third most rural county in Maine. The State of Maine is considered an "aging" state, with 17 percent of the population being 65 years old and over, a higher rate than the overall US population (14%). In Downeast PHD, 18.5% of the population is 65 years or older. The majority of the population in Downeast PHD is Caucasian (95.2%). In 2010, 2.1 percent of the population in Downeast PHD identified themselves as American Indian. Economically, the two counties within Downeast PHD differ greatly. During the period of 2008-12, the median family income in Washington was \$36,486 (lowest in the state), whereas the median income in Hancock was \$48,635 (the fifth-highest statewide). Just over 15 percent of the population in Downeast PHD lives below the poverty level. Overall, Downeast PHD tends to be older and more rural compared to the rest of the state and varies greatly in terms of socioeconomic factors.

It is within the context of these demographic characteristics that substance abuse in Downeast Public Health District (PHD) must be examined.

### Purpose of this Report

This report takes into account the primary objectives of the Office of Substance Abuse and Mental Health Services (SAMHS): to identify substance abuse patterns in defined geographical areas, establish substance abuse trends, detect emerging substances, and provide information for policy development and program planning. It also highlights all the prevention priorities identified in the SAMHS strategic plan: underage drinking, high-risk drinking, misuse of prescription drugs, and marijuana use. Finally, the report monitors many of the factors that contribute to substance use, such as access and perceptions of harm, as well as the common negative consequences such as crime, car crashes and overdose deaths.

This report includes data available through May 2015. Older and unchanged data are included when more recent data were not available. Five major types of indicators are included: self-reported substance consumption, consequences of substance use, factors contributing to substance use, indicators about mental health and substance abuse, and treatment admissions.

The most recent data available for the Behavioral Risk Factor Surveillance System (BRFSS) are from 2013. Due to methodological changes in weighting and sampling, **2011 BRFSS data cannot be trended with previous BRFSS years.** In addition, please note that **data results from the 2013 Maine Integrated Youth Health Survey are not available for the Downeast Public Health District due to an insufficient sample size.**

Previous reports are available at the [www.maineosa.org](http://www.maineosa.org) website. For additional data and resources please visit the Maine State Epidemiological Outcomes Workgroup (SEOW) data dashboard at [www.maineoseow.com](http://www.maineoseow.com).



## Consumption of Substances

---

Consuming harmful substances can have detrimental effects on an individual's well-being, including increased risks of morbidity, addiction and mortality, and has a harmful effect on society as a whole including increased motor vehicle accidents and crime. However, it is the manner and frequency with which people drink, smoke and use drugs that are often linked to particular substance-related consequences. To understand fully the magnitude of substance use consequences, it is important to first understand the prevalence of substance use consumption, itself. Consumption includes overall use of substances, acute or heavy consumption and consumption by high risk groups (e.g., youth, college students, pregnant women).

As demonstrated by the indicators below, alcohol remains the substance most often used by Downeast PHD residents across the lifespan. Alcohol and marijuana use among youth and younger adults continues to be a concern. Downeast PHD made great gains in reducing underage drinking, marijuana use, prescription drug use and tobacco use. According to the most recent data available, tobacco use appears to be slightly higher among high school students in Downeast PHD when compared to the rest of the state. Marijuana is the most commonly used drug in Maine and Downeast PHD is no different in this respect, showing the highest rate of use among the public health districts.

## Alcohol


**Indicator Description: ALCOHOL USE AMONG YOUTH.** This measure shows the percentage of Maine high school students who reported having had one or more alcoholic drinks within 30 days prior to the survey.

**Why Indicator is Important:** Alcohol is the most often used substance among youth in Maine. In addition to the risks alcohol consumption carries for adults, developing adolescent brains are especially susceptible to the health risks of alcohol consumption. Adolescents who consume alcohol are more likely to have poor grades and be at risk for experiencing social problems, depression, suicidal thoughts, assault, and violence.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** Almost one in four high school students in Downeast PHD (24%) reported having consumed one or more alcoholic beverages in the past 30 days in 2011. This was lower than the statewide average (28%).

**Figure 1. Percent of high school students by Public Health District who had at least one drink of alcohol during past 30 days: 2009-2013**



Source: MIYHS


**Indicator Description: HIGH-RISK ALCOHOL USE AMONG YOUTH.** This indicator presents the percentage of Maine high school students who reported having had five or more alcoholic drinks in a row in one sitting at least once during the 30 days prior to the survey.

**Why Indicator is Important:** Youth are more likely than adults to engage in high-risk drinking when they consume alcohol. High-risk alcohol use contributes to violence and motor vehicle crashes and can result in negative health consequences for the consumer, including injuries and chronic liver disease. Youth who engage in high-risk drinking also are more likely to use drugs and engage in risky and antisocial behavior.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, almost one in seven high school students in Downeast PHD (15%) reported having consumed five or more alcoholic beverages in one sitting during the past 30 days. This was similar to the statewide average of 15 percent; and considerably lower than in 2009.

**Figure 2. Percent of high school students by Public Health District who had at least five drinks in a row during past 30 days: 2009-2013**



Source: MIYHS




**Indicator Description: ALCOHOL USE AMONG ADULTS.** This indicator portrays the percentage of adults who reported having consumed one or more alcoholic drinks on one or more days within the past 30 days.

**Why Indicator is Important:** Alcohol is the most often used substance in Maine adults. Excessive and high-risk alcohol use may contribute to violence and result in many negative health consequences for the consumer. Moderate drinking can also have negative health effects and lead to such consequences as alcohol-related motor vehicle crashes and increased injuries. Current alcohol use in pregnant women is also linked to low birth weight babies, sudden infant death, and other developmental delays in children.

**Data Source(s):** BRFSS, 2011-2013.

**Summary:** The percentage of adults in Downeast PHD reporting drinking at least one alcoholic beverage within the past 30 days decreased from 2011 (60%) to 2013 (56%).

**Figure 3. Percent of adults by Public Health District who reported drinking during past 30 days: 2011-2013**



Source: BRFSS


**Indicator Description: AT RISK FROM HEAVY ALCOHOL USE.** This indicator examines the percentage of Maine residents who are at risk from heavy drinking in the past month. Heavy drinking is defined as two drinks per day for a man or one drink per day for a woman.

**Why Indicator is Important:** Heavy drinking increases the risk for many health and social related consequences. People who consume alcohol heavily are at increased risk for a variety of negative health consequences, including alcohol abuse and dependence, liver disease, certain cancers, pancreatitis, heart disease, and death. It has also been found that the more heavily a person drinks the greater the potential for problems at home, work, and with friends.<sup>1</sup>

**Data Source(s):** BRFSS, 2011-2013.

**Summary:** The percentage of adults in Downeast PHD at risk from heavy alcohol use (1-2 drinks per day) increased from 2012 (6%) to 2013 (10%). This rate was slightly above the statewide average (7%) and the highest among the public health districts.

**Figure 4. Percent of adults by Public Health District who reported heavy drinking during past 30 days: 2011-2013**



Source: BRFSS

<sup>1</sup> Citation from Alcoholscreening.org, a service of Join Together and the Boston University School of Public Health. Retrieved from <http://www.alcoholscreening.org/Learn-More.aspx?topicID=8&articleID=26> on 7/11/2014.


**Indicator Description: HIGH-RISK ALCOHOL USE AMONG ADULTS.** This indicator reflects the percentage of adults who reported engaging in high-risk “binge” drinking within the past 30 days. This is defined as five or more drinks in one sitting for a male and four or more drinks in one sitting for a female.

**Why Indicator is Important:** Binge drinking is considered to be a type of high-risk drinking, meaning it increases the risk for many health- and social-related consequences. It has been linked to injury (such as falls, fights, and suicides), violence, crime rates, motor vehicle crashes stroke, chronic liver disease, addiction, and some types of cancer.

**Data Source(s):** BRFSS, 2011-2013.

**Summary:** The rate of binge drinking during the past 30 days among adults in Downeast PHD decreased from 17 percent in 2012 to 15 percent in 2013. This was slightly better than the statewide average (17%).


**Figure 5. Percent of adults by Public Health District who reported binge drinking during past 30 days: 2011-2013**



Source: BRFSS

**Summary:** During 2011-13 (combined years), the highest rate of binge drinking in Downeast PHD was among the 18 to 25 year old population at 32 percent, followed by 26 to 35 year olds at 28 percent, 36 to 49 year olds (20%), and residents over the age of 50 and older (9%). Rates differed little from those of the state.

**Figure 6. Percent of adults by Public Health District who reported binge drinking in past 30 days by age group: 2011-13**



Source: BRFSS

## Tobacco


**Indicator Description: SMOKING AMONG YOUTH.** This indicator illustrates the percentage of Maine high school students who reported smoking a cigarette on at least one occasion within 30 days prior to the survey.

**Why Indicator is Important:** Use of tobacco is associated with a greater risk of negative health outcomes, including cancer, cardiovascular, and chronic respiratory diseases, as well as death.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, almost one in six high school students (17%) in Downeast PHD reported having smoked one or more cigarettes in the past 30 days. This was similar to the state average (16%) and an improvement over the 2009 rate (21%).

**Figure 7. Percent of high school students by Public Health District who reported smoking one or more cigarettes during past 30 days: 2009-2013**



Source: MIYHS


**Indicator Description: SMOKELESS TOBACCO AMONG YOUTH.** This indicator illustrates the percentage of Maine high school students who reported using smokeless tobacco on at least one occasion within 30 days prior to the survey.

**Why Indicator is Important:** Use of tobacco is associated with a greater risk of negative health outcomes, including cancer, cardiovascular, and chronic respiratory diseases, as well as death.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, the percentage of high school students in Downeast PHD who report having used smokeless tobacco in the past 30 days was on par with the statewide average (8%).

**Figure 8. Percent of high school students by Public Health District who used smokeless tobacco in the past 30 days: 2009-2013**



Source: MIYHS


**Indicator Description: SMOKING AMONG ADULTS.** This indicator illustrates the percentage of Maine adults who reported using cigarettes on at least one occasion within 30 days prior to the survey.

**Why Indicator is Important:** Smoking is associated with a greater risk of negative health outcomes, including cancer, cardiovascular and chronic respiratory diseases, as well as death.

**Data Source(s):** BRFSS, 2011-2013.

**Summary:** The percentage of adults in Downeast PHD who reported having smoked at least one cigarette within the past 30 days decreased by two percentage points from 2012 (21%) to 2013 (19%). This rate was similar to the statewide average (20%).

**Figure 9. Percent of adults by Public Health District who reported smoking a cigarette in the past 30 days: 2011-2013**



Source: BRFSS

## Prescription Drugs


**Indicator Description: MISUSE OF PRESCRIPTION DRUGS AMONG YOUTH.** This indicator presents the percentage of Maine high school students who reported using prescription drugs that were not prescribed to them by a doctor within 30 days prior to the survey.

**Why Indicator is Important:** Some people are using available prescription drugs, including stimulants and opiates, instead of illegal drugs to get high. Abuse of prescription drugs may lead to consequences such as unintentional poisonings or overdose, automobile crashes, addiction, and increased crime.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, five percent of high school students in Downeast PHD reported having taken prescription drugs not prescribed to them by a doctor one or more times in the past 30 days, compared to seven percent reporting similar activity statewide.

**Figure 10. Percent of high school students by Public Health District who have taken prescription drugs not prescribed to them by a doctor (past 30 days): 2009-2013**



Source: MIYHS




**Indicator Description: MISUSE OF PRESCRIPTION DRUGS AMONG ADULTS.** This measure reflects the percentage of adults in Maine who reported using prescription drugs not prescribed to them by a doctor, or using them in a way other than the one prescribed, at least once in their lifetime. Because of small sample sizes, survey data from multiple years must be combined in order to produce this estimate.

**Why Indicator is Important:** Some Mainers misuse available prescription drugs (including stimulants and opiates) instead of illegal drugs to get high. Abuse of prescription drugs may lead to consequences such as unintentional poisonings, overdose, dependence and increased crime.

**Data Source(s):** BRFSS, 2011-13

**Summary:** During 2011-13 (combined years), over three percent (3.6%) of adults in Downeast PHD reported to have misused prescription drugs in their lifetime. This was slightly above the statewide average (3.3%).

**Figure 11. Misuse of prescription drugs among Maine residents (18 and older) in their lifetime, by gender and public health district: 2011-13**



Source: BRFSS


**Indicator Description: NONMEDICAL USE OF PRESCRIPTION PAIN RELIEVERS AMONG MAINERS AGE 18 AND OLDER.** This measure reflects the percentage of adults who reported using prescription pain relievers in the past year, for reasons other than their intended purpose. Because of small sample sizes, survey data from multiple years must be combined in order to produce this estimate.

**Why Indicator is Important:** Mainers are increasingly using available prescription drugs, particularly pain relievers, instead of illegal drugs to get high. Abuse of prescription drugs may lead to consequences such as unintentional poisonings, overdose, dependence and increased crime.

**Data Source(s):** NSDUH, 2010-12.

**Summary:** In 2010-12, 10.6 percent of 18 to 25 year olds in Aroostook/Downeast reported non-medical use of prescription pain relievers during the past year, compared to 2.9 percent among those who were 26 and older.


**Figure 12. Percent of population 18 years old or older who used prescription pain relievers in past year for nonmedical use, by Public Health District: 2010-12**



Source: NSDUH

**Summary:** Past year use of non-medical pain relievers among 18 to 25 year olds in Aroostook/Downeast decreased by two percentage points from 2004-06 (12.7%) to 2010-12 (10.6%). This was on par with rates at the state level.

**Figure 13. Percent of 18 to 25 year olds who used prescription pain relievers in past year for nonmedical use, by Public Health District: 2004-06 to 2010-12**



Source: NSDUH

## Other Illegal Drugs


**Indicator Description: CURRENT MARIJUANA USE.** This measure shows the percentage of Maine residents who reported using marijuana in the past 30 days. This is presented for high school students and adults in Maine.

**Why Indicator is Important:** Marijuana can be addictive and is associated with increased risk for respiratory illnesses and memory impairment. Even occasional use can have consequences on learning and memory, muscle coordination, and mental health symptoms.

**Data Source(s):** MIYHS, 2009-2013; BRFSS, 2012; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, 20 percent (one in five) of high school students in Downeast PHD reported having used marijuana one or more times in the past 30 days compared to 22 percent statewide.


**Figure 14. Percent of high school students by Public Health District who have used marijuana during past 30 days: 2009-2013**



Source: MIYHS

**Summary:** Among Downeast PHD adults, 8.1 percent reported using any marijuana within the past 30 days. This was slightly higher than the statewide rate of 7.4 percent. Rates in Downeast PHD were higher among males (10.7%) as compared to females (6.1%).

**Figure 15. Percent of adults who have used marijuana during the past 30 days, by gender and Public Health District: 2012-13**



Source: BRFSS


**Indicator Description: LIFETIME COCAINE USE AMONG YOUTH.** This indicator illustrates the percentage of Maine high school students who used cocaine at least once in their lifetime (i.e., ever).

**Why Indicator is Important:** Cocaine is highly addictive. Use of cocaine is associated with adverse health effects such as cardiac events, seizures, and stroke. It also increases the risk of cognitive impairment, injury, and crime.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, six percent of high school students in Downeast PHD reported that they had used cocaine (in any form) during their lifetime; this was similar to the statewide average of seven percent.

**Figure 16. Percent of high school students by Public Health District that have used cocaine in any form during their lifetime: 2009-2013**



Source: MIYHS

## Consequences Resulting from Substance Use and Abuse

---

Both individuals and communities suffer the consequences of substance abuse in terms of increased health care needs and criminal justice resources. While a great deal of information regarding substance use can be obtained from the data described in the previous section, information on the effects of that use on individuals and communities can be derived from what has come to be called “consequence” data. Consequences are defined as the social, economic and health problems associated with the use of alcohol and illicit drugs. Examples are things such as illnesses related to alcohol, drug overdose deaths, property and personal crimes, as well as driving accidents, poisonings and suicides that involve alcohol or drugs.

Although the overall rate of outpatient admissions due to substance abuse in Downeast PHD is lower than the statewide rate, the district has the highest rate of outpatient admissions due to opiates. Similarly, the rate of overdose emergency responses is lower than the state, but those that do occur are more likely to be medication related. Drug offense arrests made the Maine Drug Enforcement Agency tend to be related to pharmaceutical narcotics rather than other types of drugs.

## Substance Use and Pregnancy

---

**Indicator Description: BABIES BORN AFFECTED BY SUBSTANCES.** This measure reflects the number of infants born in Maine where a healthcare provider reported to OCFS that there was reasonable cause to suspect the baby may be affected by drugs or alcohol or demonstrating withdrawal symptoms resulting from prenatal drug exposure (illicit or prescribed) or who have fetal alcohol spectrum disorders. This measure potentially excludes instances where the infant was exposed to substances and did not show withdrawal symptoms after birth, instances where the birth of an infant affected by substances was not reported to OCFS, and any other instances in which there were discrepancies between reporters when interpreting Title 22, §4011-A; notification of prenatal exposure to drugs or having fetal alcohol spectrum disorders.


**Why Indicator is Important:** Prenatal exposure to alcohol, tobacco, and illicit drugs has the potential to cause a wide spectrum of physical, emotional, and developmental problems for these infants. The harm caused to the child can be significant and long-lasting, especially if the exposure is not detected and the effects are not treated as soon as possible.

**Data Source(s):** OCFS/MACWIS, 2010-2014

**Summary:** The rate of drug affected babies in Downeast PHD steadily increased from 2009 (3.9 per 10,000 residents) to 2013 (8.8 per 10,000 residents). In 2013, Downeast PHD had the fourth highest rate among public health districts. In 2013, rates among public health districts ranged from 2.8 drug affected babies per 10,000 residents in York to 11.6 per 10,000 residents in Penquis.



Figure 17. Number of drug affected baby notifications, by Public Health District: 2010-2013



Source: OCFS/MACWIS

## Referral Services

---

**Indicator Description: INFORMATION CALLS FOR REFERRAL SERVICES.**


2-1-1 Maine is a telephone and internet service that provides information and referrals to a wide range of health and human services. This indicator reflects the number of calls received by Maine 2-1-1 where the callers were seeking services related to the treatment of substance use. Callers are referred to support services such as alcohol anonymous meetings, residential treatment programs, outpatient counseling/certification programs, and medication assisted treatment programs. Virtually all callers are seeking treatment for themselves, family members, friends, or significant others.

**Why Indicator is Important:** The data collected provides valuable information serving as a barometer of health and human service needs related to substance use in the state.

**Data Source(s):** 2-1-1 Maine, 2010-2014

**Summary:** In 2104, Downeast PHD observed just over 10 2-1-1 Maine calls per 10,000 residents related to substance use. From 2010 to 2014, Downeast PHD held the second lowest rates among public health districts. Downeast PHD's rates have remained relatively unchanged over the past several years.

**Figure 18. Number of 2-1-1 referral calls relating to substance use per 10,000 residents, by Public Health District: 2010-2013**



Source: 2-1-1 Maine

## Criminal Justice Involvement

---

**Indicator Description: ANNUAL VIOLENT CRIME RATE.** This indicator shows the number of violent crimes reported to the police, per 10,000 people. Violent crimes include murder, rape, robberies, and aggravated assaults. The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.


Operationalized as:  $\left( \frac{\# \text{ of violent crimes}}{\text{population}} \right) \times 10,000$

**Why Indicator is Important:** Violence is associated with alcohol, though the causal pathway is not completely understood. Drinking on the part of the victim or a perpetrator can increase the risk of assaults and assault-related injuries. Estimates have indicated that at least 23 percent of sexual assaults and 30 percent of physical assaults can be attributed to alcohol. Reported violent crimes are an under-report of the total number of actual violent crimes.

**Data Source(s):** DPS, UCR, 2009-2013.

**Summary:** In 2013, there were 10.1 violent crimes per 10,000 people in Downeast PHD, compared to 12.5 per 10,000 people statewide. Although not shown in the chart, the number of violent crimes in Downeast PHD decreased from 102 in 2011 to 87 in 2013; representing a decrease of 15 percent.

Figure 19. Violent crime rate per 10,000, by Public Health District: 2009-2013



Source: DPS; UCR

**Indicator Description: ANNUAL ALCOHOL-RELATED ARREST RATE.** This indicator reflects arrests related to alcohol per 10,000 people. Alcohol-related arrests include Operating Under the Influence (OUI), liquor law violations, and drunkenness. The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.


Operationalized as:  $\left( \frac{\# \text{ of alcohol arrests}}{\text{population}} \right) \times 10,000$

**Why Indicator is Important:** OUI and liquor law arrest rates can be an indication of the rate of criminal behavior, but it is important to note that they are also an *indication of the level of law enforcement*. Arrest rates are expected to increase with increased enforcement regardless of whether a decline in criminal behavior is observed. The educational component of Maine's Driver Education and Evaluation Program serviced 5,192 Maine residents during the 2014 fiscal year.

**Data Source(s):** DPS, UCR, 2009-2013.

**Summary:** After observing a spike in 2012 (72.6 per 10,000), Downeast PHD's alcohol-related arrest rate dropped to 64.4 per 10,000 residents in 2013. The statewide alcohol-related arrest rate was 71.2 per 10,000 residents in 2013. Downeast PHD's rate has fluctuated over the past several years, ranging between 65 and 85 alcohol-related arrests per 10,000 residents.

**Figure 20. Alcohol-related arrest rate per 10,000, by Public Health District:  
2009-2013**



Source: DPS; UCR

**Indicator Description: ANNUAL DRUG-RELATED ARREST RATE.** This indicator reflects the number of arrests (made by all local and state law enforcement) that were related to drugs per 10,000 people. Drug-related arrests include manufacturing, sales, and possession. The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time as well as make relative comparisons between small and large population areas.


Operationalized as:  $\left( \frac{\# \text{ of drug arrests}}{\text{population}} \right) \times 10,000$

**Why Indicator is Important:** Arrest rates for drug sales, manufacturing and drug possession can be an indication of the rate of criminal behavior, but it is important to note that they are also an *indication of the level of law enforcement*. Arrests rates are expected to increase with increased enforcement regardless of whether a decline in criminal behavior is observed.

**Data Source(s):** DPS, UCR, 2009-2013.

**Summary:** In 2013, there were 30.3 drug-related arrests per 10,000 people in Downeast PHD; this was lower than the statewide rate (42.1 per 10,000) and the lowest among public health districts.

**Figure 21. Drug-related arrest rate per 10,000, by Public Health District: 2009-2013**




Source: DPS; UCR



**Summary:** In 2013, most drug related arrests by local and state law enforcement were related to marijuana (12.4 per 10,000 residents) followed by other dangerous non-narcotics (8.6 per 10,000), synthetic narcotics (6.1 per 10,000), and opium, cocaine, and derivatives (3.2 per 10,000).

**Figure 22. Drug-related arrest rate per 10,000, by drug type and Public Health District: 2013**



Source: DPS; UCR

\*Derivatives include cocaine/crack, codeine, heroin, and morphine.

\*\*Other dangerous non-narcotics include but are not limited to benzodiazepines, steroids, stimulants, synthetic cannabis, bath salts, methamphetamine, hallucinogens, and barbiturates.


**Indicator Description: MAINE DEA DRUG OFFENSE ARRESTS BY TYPE.** This indicator reflects drug offense arrests made by the Maine’s Drug Enforcement Agency, overall and by drug type. The MDEA, through its eight regional multi-jurisdictional task forces, is the lead state agency in confronting drug trafficking crime. This indicator differs from the previous drug-related arrest data in that it only tracks MDEA efforts and does not encompass all activity within Maine law enforcement agencies.

**Why Indicator is Important:** Drug offense arrest rates can be an indication of the rate of criminal behavior, but it is important to note that they are also an indication of the level of law enforcement. Drug arrest rates are expected to increase with increased enforcement regardless of whether a decline in criminal behavior is observed.

**Data Source(s):** MDEA-UCR, 2010-2014

**Summary:** In 2014, there were 6.2 drug offense arrests per 10,000 residents in Downeast PHD; the third highest rate among public health districts in Maine. During 2011-14 (combined years), most DEA drug offense arrests in Downeast PHD were related to pharmaceutical narcotics (3.2 arrests per 10,000) followed by Heroin (1.4 arrests per 10,000).


**Figure 23. Maine DEA Drug offense arrests per 10,000 residents, by Public Health District: 2010-2014**



Source: MDEA-UCR

**Summary:** During 2012-14 (combined years), most drug offense arrests in Downeast PHD were related to pharmaceutical narcotics (3.2 arrests per 10,000) followed by Heroin (1.4 arrests per 10,000). Although not shown, during the 2012-14 period, 17 percent (84 arrests) of pharmaceutical narcotics related arrests occurred in Downeast PHD.

**Figure 24. Maine DEA Drug offense arrests per 10,000 residents, by Public Health District: 2012-14**



Source: MDEA-UCR

\*Synthetic cathinones are synthetic derivatives of an alkaloid that are used as drugs for their stimulating properties.

## Driving Under the Influence


**Indicator Description: DRINKING AND DRIVING AMONG YOUTH.** This measure shows the proportion of high school students who reported that they drove a car after consuming alcohol at least once within 30 days prior to taking the survey.

**Why Indicator is Important:** Operating a vehicle after consuming alcohol increases the risk of motor vehicle crashes, injuries and death.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, approximately four percent of high school students in Downeast PHD reported driving a vehicle at least once after drinking alcohol in the past 30 days compared to five percent statewide reporting such behavior.

**Figure 25. Percent of high school students by Public Health District who reported drinking and driving during the past 30 days: 2009-2013**



Source: MIYHS


**Indicator Description: YOUTH AS PASSENGERS IN VEHICLES DRIVEN BY INDIVIDUALS USING ILLEGAL DRUGS.** This measure shows the proportion of high school students who reported that within 30 days prior to taking the survey they were a passenger in a car being operated by an individual who had consumed illegal drugs.

**Why Indicator is Important:** Operating a vehicle while under the influence of drugs increases the risk of motor vehicle crashes, injuries and death.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size. .

**Summary:** In 2011, 18 percent of high school students in Downeast PHD reported that, within the past 30 days, they had been passengers in a vehicle operated by someone who had taken illegal drugs compared to 22 percent across the state. This represents a notable decrease in the rate of such incidents from 2009 to 2011.

**Figure 26. Percent of high school students by Public Health District who rode in a vehicle driven by someone who had taken illegal drugs: 2009-2013**



Source: MIYHS

**Indicator Description: ALCOHOL/DRUG-INVOLVED MOTOR VEHICLE CRASH RATE.** This indicator shows the number of motor vehicle crashes in which alcohol or drugs were a factor per 10,000 people. Due to new data collection regulations, crash rate data is no longer separated by alcohol and drugs. Alcohol and drugs are now combined into one rate. Alcohol/drug-involved crashes means that at least one driver had consumed alcohol or drugs prior to the crash. The rate per 10,000 allows us to see the frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.

Operationalized as:  $\left( \frac{\# \text{ of alcohol/drug-involved crashes}}{\text{population}} \right) \times 10,000$

**Why Indicator is Important:** Motor vehicle crashes are the second-leading cause of traumatic brain injury, with 27 percent of traumatic brain injuries occurring from motor vehicle crashes.<sup>2</sup> In 2009, alcohol was attributed to 96 percent of the alcohol/drug-related crashes statewide.


**Data Source(s):** MDOT/MBHS, 2010-2014.

**Summary:** Although Downeast PHD still has one of the highest alcohol/drug involved crash rates among public health districts, the rate has decreased from 2012 (13.1 crashes per 10,000 residents) to 2014 (9.8 crashes per 10,000 residents).

---

<sup>2</sup> 2007 Maine Injury Report, Maine Center for Disease Control, Injury Prevention Program. Retrieved 7/11/2014 from <http://www.maine.gov/dhhs/mecdc/population-health/inj/documents/2007maineinjuryreport.pdf>

**Figure 27. Alcohol/Drug-related motor vehicle crash rate per 10,000, by  
Public Health District: 2010-2014**



Source: MDOT/MBHS

## Hospital Visits Related to Substance Use

---

**Indicator Description: INPATIENT ADMISSIONS RELATED TO SUBSTANCE USE.** This indicator shows the number of inpatient hospital admissions (per 10,000 people) where alcohol, opiates, or other drugs were recorded as the primary diagnosis for which services were sought at admission. “Inpatient” refers to a patient whose treatment needs at least one night's residence in a hospital. The substance for which treatment was received was identified through hospital codes (ICD-9 codes) and includes those related to alcohol and psychoactive substances (303-305). More than one substance may be involved in a single visit. The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.

Operationalized as:  $\left( \frac{\# \text{ of inpatient hospitalizations}}{\text{population}} \right) \times 10,000$


**Why Indicator is Important:** Hospital admissions related to substance use are an indication of injury sustained through substance use and the impact it has on the healthcare system.

**Data Source(s):** MHDO, 2010 and 2011.

**Summary:** The inpatient admissions rate due to substance abuse in Downeast PHD changed very little from 2010 (79.3 admissions per 10,000 residents) to 2011 (80.8 admissions per 10,000). In 2011, Downeast PHD had a lower inpatient rate than the state (95.3 admissions per 10,000) and had the third lowest rate among public health districts.



**Figure 28. Inpatient hospital admissions (per 10,000 people) related to substance use\*, by Public Health District: 2010, 2011**




Source: MHDO

\*Visits may involve multiple substances

**Summary:** In 2011, most inpatient admissions due to substance use in Downeast PHD were related to alcohol (53.3 admissions per 10,000), followed by opiates (21.2 admissions per 10,000), marijuana (10.5 admissions per 10,000), and other drugs (12.8 admissions per 10,000). Among public health districts, Downeast PHD tends to have the third lowest inpatient rate across all the substances.

**Figure 29. Inpatient hospital admissions (per 10,000 people) related to substance use\*, by Public Health District and drug type: 2011**



Source: MHDO

\*Visits may involve multiple substances

**Indicator Description: OUTPATIENT HOSPITAL VISITS RELATED TO SUBSTANCE USE.** This indicator shows the of outpatient hospital admissions (per 10,000 people) where alcohol, opiates, or other drugs was recorded as the primary diagnosis for which services were received. “Outpatient” refers to patients who receive treatment at a hospital or clinic but are not admitted overnight. The substance for which treatment was received was identified through hospital codes (ICD-9 codes) and includes those related to alcohol psychoactive substances (303-305). The rate per 10,000 allows us to see frequency with which an occurrence shows up within a population over time, as well as make relative comparisons between small and large population areas.


Operationalized as:  $\left( \frac{\# \text{ of outpatient hospitalizations}}{\text{population}} \right) \times 10,000$

**Why Indicator is Important:** Outpatient hospital visits related to substance use are an indication of injury sustained through substance use and the impact it has on the healthcare system.

**Data Source(s):** MHDO, 2010 and 2011

**Summary:** From 2010 to 2011, the outpatient admission rate to due to substance use in Downeast PHD decreased from 973.3 admissions per 10,000 residents to 681.6 admissions per 10,000 residents. This was lower than the statewide rate in 2011.

**Figure 30. Outpatient hospital admissions (per 10,000 people) related to substance use,\* by Public Health District: 2010, 2011**




Source: MHDO

\*Visits may involve multiple substances

**Summary:** In 2011, Downeast had the highest outpatient admission rate due to opiates among public health districts (434.7 admissions per 10,000); the statewide rate was 300.4 admissions per 10,000. The rate of outpatient hospital admissions due to alcohol in Downeast PHD s was 176.9 per 10,000 residents which was lower than the state and third lowest among the public health districts.

**Figure 31. Outpatient hospital admissions (per 10,000 people) related to substance use\*, by Public Health District and drug type: 2011**



Source: MHDO

\*Visits may involve multiple substances

\*\*Includes prescription narcotics, methadone, and heroin.


**Indicator Description: POISONING CASES DOCUMENTED BY THE POISON CENTER.** This measure reflects the rate of calls to the Northern New England Poison Center in which the Center determined that a poisoning occurred. These calls are for the state of Maine only. The Center reports poisonings in three categories: unintentional, meaning those that are accidental; suspected substance abuse cases, meaning cases where the Center believes the intent is for an individual to get high; and suspected suicides, meaning staff at the Center determine that the individual attempted suicide. The categories reflect the caller's self-report and are not considered clinical or medical diagnoses.

**Why Indicator is Important:** The exposure to and ingestion of damaging substances can have many physiologic side effects. Poisonings can be influenced by programs to prevent substance abuse, accidental poisoning, suicide and fatal interaction among medications.

**Data Source(s):** NNEPC, 2012-14

**Summary:** During the period of 2012-14 the Poison Center received 10.0 calls per 10,000 residents in Downeast PHD that were suspected to be attempted suicide; this was lower than the statewide rate (12 calls per 10,000 residents). Within the same period, Downeast PHD observed 2.9 calls per 10,000 residents that were suspected to be substance abuse; this was also on par with the statewide rate of 3.4 calls per 10,000 residents.


**Figure 32. Number of poisonings reported to New England Poison Center 10,000 residents, by intent and Public Health District: 2012-14**



Source: NNEPC


**Summary:** Downeast PHD's rate for calls suspected to be substance abuse increased from 2.0 calls per 10,000 residents in 2008-10 to 2.9 calls per 10,000 in 2012-14. As for calls suspected to be suicide attempts, Downeast PHD's rate remained 10.0 calls per 10,000 residents in 2008-10 and 2012-14.

**Figure 33. Number of poisonings reported to New England Poison Center suspected to be substance abuse per 10,000 residents, by Public Health District: 2012-14**





**Figure 34. Number of poisonings reported to New England Poison Center suspected to be suicide attempts per 10,000 residents, by Public Health District: 2012-14**



## Overdoses and Related Deaths

---

**Indicator Description: OVERDOSES.** This indicator shows the rate of persons receiving help from Emergency Medical Services (EMS) related to an overdose.


**Why Indicator is Important:** Overdosing on a substance can cause serious physical harm resulting in hospitalization and even death. Responding to overdoses also uses valuable EMS resources. The rate per 10,000 allows us to see the frequency with which an occurrence happens within a population over time, as well as make relative comparisons between small and large population areas. In this case, the base of 10,000 people was used due to small numbers.

Operationalized as:  $\left( \frac{\# \text{ of overdose deaths}}{\text{population}} \right) \times 10,000$

**Data Source(s):** EMS, 2011-2014


**Summary:** In 2014, Downeast PHD had a low rate of drug/medication overdose responses among public health districts with 17.3 per 10,000 residents. This was compared to the statewide rate of 22.2 drug/medication responses per 10,000 residents. As for overdose responses related to alcohol, Downeast PHD held a rate of 15.8 per 10,000 residents; this was slightly lower than the state average (16.6 per 10,000). Downeast PHD's rate of overdose responses classified as drug/medication and alcohol related has remained relatively stable since 2011.

Figure 35. Number of overdose EMS responses per 10,000 residents, by Public Health District and primary type of substance involved: 2014




Source: Emergency Medical Services, 2012

Figure 36. Number of overdose EMS responses due to drug/medication per 10,000 residents, by Public Health District: 2011-2014



Source: Emergency Medical Services, 2012

**Figure 37. Number of total overdose EMS responses due to alcohol per 10,000 residents, by Public Health District: 2011-2014**



Source: Emergency Medical Services


**Indicator Description: NALOXONE ADMINISTRATIONS.** This indicator shows the number of persons receiving naloxone administrations from Emergency Medical Services (EMS) related to an opioid overdose. Naloxone is a medication administered to patients who have experienced an overdose related to an opioid (e.g., prescription painkillers, heroin, and morphine). This indicator includes instances where the opioid overdose is accidental (that is, not a result of intentional or recreational misuse).

**Why Indicator is Important:** Overdosing on a substance can cause serious physical harm resulting in hospitalization and even death. Responding to overdoses also uses valuable EMS resources. Furthermore, it is worth stating that this indicator gives us a better sense of the prevalence of opioid overdoses since it includes those that did not result in death.

**Data Source(s):** Emergency Medical Services, 2014

**Summary:** In 2014, Downeast PHD observed a rate of 4.4 naloxone administrations per 10,000 residents via overdose ambulance responses; this was below the statewide rate of 6.2 naloxone administrations per 10,000 residents. Although not shown, naloxone was administered by EMS responders a total of 38 times in 2014.

**Figure 38. Naloxone\* administrations per 10,000 residents, by Public Health District: 2014**



Source: Emergency Medical Services

\*Naloxone is a medication administered to counter the effects of an overdose due to opioids.

**Indicator Description: DEATHS DUE TO OVERDOSE.** This measure reflects the number of deaths where the cause of death was directly related to the consumption of one or more substances. The measure excludes deaths where a substance may have been ingested prior to engaging in a behavior that resulted in death (e.g., drunk driving) or where lifetime substance use and abuse may have impacted health (e.g., cirrhosis). In order to preserve anonymity and strengthen validity, rates were calculated based on the sum of deaths per three year intervals. The rate per 100,000 allows us to see the frequency with which an occurrence happens within a population over time, as well as make relative comparisons between small and large population areas. In this case, the base of 100,000 people was used due to small numbers.


Operationalized as:  $\left( \frac{\# \text{ of overdose deaths}}{\text{population}} \right) \times 100,000$

**Why Indicator is Important:** One of the most extreme consequences of alcohol and drug abuse is overdose death; that is, the substance(s) consumed played a direct role in an individual's death. These are seen as potentially preventable deaths.

**Data Source(s):** Marci Sorg, Margaret Chase Smith Policy Center at University of Maine, Office of the Chief Medical Examiner, 2009-10 to 2013-14.

**Summary:** During 2013-14 (combined years), Downeast PHD observed an average of 12.2 drug related overdose deaths per 100,000 residents; this was similar to the statewide average of 12.9 drug related overdose deaths per 100,000 residents.

**Figure 39. Drug-related death rate per 100,000, by Public Health District:  
2009-10 to 2013-14**



Source: Marci Sorg, Margaret Chase Smith Policy Center at University of Maine, Office of the Chief Medical Examiner



## Factors Contributing to Substance Use and Abuse

---

A body of substance abuse prevention research has identified certain groups of factors that “cause” or have an impact on substance use and the consequences related to use. That is, they appear to influence the occurrence and magnitude of substance use and its related consequences. Generically, these causal factors (also known as contributing factors) are categorized into groups which include:

- Social Access (e.g., getting drugs and alcohol from friends or family)
- Retail Availability (e.g., retailer not carding properly)
- Pricing & Promotion (e.g., two-for-one specials, industry sponsorships or signage)
- Social/Community Norms (e.g., parental/community attitudes and beliefs)
- Enforcement (e.g., lack of compliance checks)
- Perceptions of Harm (e.g., individuals’ belief that using a substance is harmful)<sup>3</sup>
- Perceived Risk of Being Caught (e.g., individuals’ belief that s/he will be caught by parents or police)<sup>4</sup>

Substance abuse prevention in Maine is undertaken with the assumption that making changes to these factors at the community level will result in changing behaviors around substance use and related problems. It is through positively impacting these factors that Maine can achieve population-level changes in substance consumption and consequences.

Although most high school students in Downeast PHD seem to perceive that regular use of substances poses a risk of harm, many do not think they will be caught by their parents or police if they use alcohol or marijuana. According to the most recent available data, most students in Downeast PHD think it is easy to obtain alcohol and marijuana. In 2014, Downeast PHD had the highest rate of liquor licensees per capita than any other public health district.

---

<sup>3</sup> Bonnie, R. J & O’Connell, M.E., Eds. (2004). *Reducing Underage Drinking: A Collective Responsibility*. The National Academies Press: Washington, DC.

<sup>4</sup> Birckmayer, J. D., Holder, H. D., Yacoubian, Jr., G. S., & Friend, K. B. (2004). A general causal model to guide alcohol, tobacco, and illicit drug prevention: assessing the research evidence. *Journal of Drug Education*, 34(2), 121-153.

## Availability and Accessibility


**Indicator Description: PERCEIVED EASE OF OBTAINING ALCOHOL BY UNDERAGE YOUTH.** This indicator reflects the percentage of high school students (grades 9 to 12) who reported that it would be easy or very easy for them to get alcohol if they wanted some.

**Why Indicator is Important:** In 2013, students who reported that they thought alcohol was easy to obtain were three times as likely to report consuming alcohol within the past month compared to students who did not think it was easy to obtain.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** Sixty-five percent of high school students in Downeast PHD (almost two out of three) indicated that it was easy to get alcohol, compared to the statewide rate of 67 percent.

**Figure 40. Percent of high school students by Public Health District who reported it was easy to get alcohol: 2009-2013**



Source: MIYHS


**Indicator Description: PARENT PERCEPTION OF ACCESSIBILITY OF ALCOHOL AT HOME.** This indicator measures the percentage of parents reporting that their teen would be able to access alcohol they had purchased without their knowledge. This data comes from the Maine Parent Survey administered by Pan Atlantic for the Maine Office of Substance Abuse and Mental Health Services.

**Why Indicator is Important:** Easy access to alcohol at home is a major contributing factor to underage drinking.

**Data Source(s):** Parent Survey 2013 and 2015

**Summary:** In 2015, among parents of middle and high school youth, 56 percent felt it was possible for their children to access alcohol they had purchased without their knowledge; this was 6 percentage points higher than the statewide average. Downeast PHD's rate increased by 14 percentage points since 2013.

**Figure 41. Parent perception of accessibility of parent purchased alcohol without parental knowledge: 2013 and 2015**



Source: Parent Survey


**Indicator Description: PARENT PERCEPTION OF ACCESSIBILITY OF RX DRUGS AT HOME.** This indicator measures the percentage of parents reporting that their teen would be able to access prescription medication (not prescribed to their child) without their knowledge. This question was first asked in 2015 Parent Survey. This data comes from the Maine Parent Survey administered by Pan Atlantic for the Maine Office of Substance of Abuse and Mental Health Services.

**Why Indicator is Important:** Easy access to prescription drugs at home is a major contributing factor to prescription drug misuse.

**Data Source(s):** Parent Survey 2015

**Summary:** Almost a third (33%) of Downeast PHD parents felt that, at home, their child would be able to access prescription medications that were not prescribed to the child, without their parents' knowledge. Downeast PHD's rate was two percentage points lower than the statewide average.

**Figure 42. Parent perception of teen accessibility of prescription drugs at home without parental knowledge: 2015**



Source: Parent Survey


**Indicator Description: PERCEIVED EASE OF OBTAINING MARIJUANA BY YOUTH.** This indicator illustrates the percentage of high school students reporting it would be easy or very easy to obtain marijuana if they wanted it.

**Why Indicator is Important:** In 2013, students who reported that they thought marijuana was easy to obtain were nearly eight times as likely to use marijuana in the past 30 days compared to their peers who thought it was difficult to obtain.

**Data Source(s):** MIYHS, 2009-2013.

**Summary:** In 2011, more than one-half (56%) of high school students in Downeast PHD indicated that it would be easy to get marijuana; this was similar to the statewide average (58%).

**Figure 43. Percent of high school students by Public Health District who reported it would be easy to get marijuana: 2009-2013**



Source: MIYHS


**Indicator Description: NUMBER OF ALCOHOL OUTLETS PER CAPITA.** This indicator reflects the number of active (as of June 2015) retail establishments selling alcohol per person. This includes both on-premise (e.g., bars, restaurants) and off-premise (e.g., convenience stores) establishments. It is calculated by dividing the number of retail establishments by the number of residents in the county (based on 2014 U.S. Census estimates).

**Why Indicator is Important:** National research shows that there is a correlation between the number of places that sell alcohol in an area (retail density) and the rate of alcohol-related crime.<sup>5</sup>

**Data Source(s):** DPS, Liquor Licensing and Compliance, 2015; U.S. Census, 2014.

**Summary:** In 2015, the number of active liquor licenses in Downeast PHD per 1,000 residents (4.3) was the highest among all public health districts. Although not shown, as of June of 2015, Downeast PHD had 368 active liquor licensees.

**Figure 44. Number of liquor licensees per 1,000 residents, by Public Health District: 2015**



Source: DPS/U.S. Census

<sup>5</sup> Grube, J. W., Gruenewald, P. J. & Chen, M. J. (2010). Community alcohol outlet density and underage drinking. *Addiction*, 105, 270-278.


**Indicator Description: DISPENSED QUANTITY OF SCHEDULE II DRUGS PER CAPITA.** These indicators reflect the dispensed quantity of narcotics, tranquilizers, and stimulants through prescriptions in Maine. This includes only prescription drugs that are classified “Schedule II” drugs, meaning those with a higher potential for abuse and addiction. It is important to note that the dispensed quantity does not indicate the size or dosage of the pills associated with the prescription. All pharmacies in Maine report to the Prescription Monitoring Program.

**Why Indicator is Important:** The dispensed quantity per capita indicates the volume of prescription drugs potentially available in the community for diversion (e.g., gift, sale, or theft). A higher level of availability contributes to misuse by individuals without a prescription.

**Data Source(s):** PMP, 2010-2014

**Summary:** Since 2009, Downeast PHD’s dispensed quantity of narcotics per person has been gradually decreasing over the past several years. In 2014, Downeast PHD had a quantity dispensed rate of 62.4 pills per person; this was very similar to the statewide rate (60.4 pills per person). In 2014, as with all public health districts, Downeast PHD observed an uptick in the rate of narcotics dispensed per capita. This increase was most likely due to the changes in data collection and drug classification (see note below chart).

Figure 45. Dispensed quantity of narcotics per capita, by  
Public Health District: 2010-2014\*




Source: PMP, 2010-2014

\*The increase observed in 2014 in number of narcotic pills prescribed per capita was due in part to the inclusion of the previously unscheduled drug Tramadol (became a schedule II drug as of 8/18/2014) as well as the inclusion of data submitted via the Veterans Administration (as of 10/31/2014). Data shown only includes two months of VA data therefore rates are expected to increase in coming years.



**Summary:** In 2014, the rate of the quantity of stimulants dispensed in Downeast PHD was 11.6 pills per person: this was the third lowest rate among public health districts. For the past several years Downeast PHD's rate for dispensed stimulant pills per capita has remained relatively stable.


**Figure 46. Dispensed quantity of stimulants per capita, by Public Health District: 2010-2014**



Source: PMP, 2010-2014

**Summary:** In 2014, the rate of the quantity of tranquilizers dispensed in Downeast PHD was 27.8 pills per person; this was on par with the statewide average (26.7 pills per person). As with all other public health districts, the dispensed quantity rate for tranquilizers has decreased slightly since 2012.

**Figure 47. Dispensed quantity of tranquilizers per capita, by Public Health District: 2010-2014**



Source: PMP, 2010-2014


**Indicator Description: SUBSTANCES REQUESTED FOR VERIFICATION.** This indicator shows the rate of requests by non-law enforcement for medication verification through the Northern New England Poison Center. A person may call the NNEPC for many reasons, one being to help identify a medication or substance which another person has consumed or that has been found. The calls reflected in this indicator have been characterized by NNEPC as likely related to substance abuse, although NNEPC staffs do not make a formal or clinical assessment.

**Why Indicator is Important:** The increased volume of medication verification calls suggests a greater availability of those drugs in the community. This measure also suggests that there is a higher awareness among the community and parents for potential misuse of prescription pills which is prompting calls.

**Data Source(s):** NNEPC, 2013-14

**Summary:** During the period of 2013-14, most calls to NNEPC within Downeast PHD requesting substance verification involved opioids (54.3 calls per 10,000 residents), followed by benzodiazepines (30.5 calls per 10,000 residents), and stimulants (9.8 calls per 10,000 residents). These rates were mostly above statewide averages. Although not shown, calls throughout the state requesting substance verification have been decreasing steadily within all drug types across public health districts for the past several years.

**Figure 48. Number of poisonings reported to New England Poison Center per 10,000 residents, by drug type and Public Health District: 2013-14**



Source: NNEPC

## Perceived Risk and Harm


**Indicator Description: PERCEIVED RISK FROM BINGE DRINKING AMONG YOUTH.** This indicator reflects the percentage of individuals who perceive that there is moderate-to-great risk from drinking five or more drinks once or twice per week.

**Why Indicator is Important:** In 2013, High school students who did not perceive a moderate to great risk of harm from binge drinking once or twice a week were more than twice as likely to drink in the past month as high school students who did perceive risk of harm. Perceptions around the risks of binge drinking are related to high-risk alcohol use among adults as well.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, 77 percent of high school students in Downeast PHD indicated that there is a moderate-to-great risk of people harming themselves if they consume five or more drinks regularly. This was very similar to the state average (78%).

**Figure 49. Percent of high school students by Public Health District who reported a risk of harm from consuming five or more drinks once or twice per week: 2009-2013**



Source: MIYHS


**Indicator Description: PERCEIVED RISK FROM BINGE DRINKING AMONG MAINERS.** This indicator reflects the percentage of Mainers age 18 and older who perceive that there is risk from consuming five or more drinks once or twice per week. Because of small sample sizes, survey data from multiple years must be combined in order to produce this estimate.

**Why Indicator is Important:** The perception that consuming a lot of alcohol is risky indicates an individual is knowledgeable about health risks and other negative consequences. Adults are less likely to binge drink if they perceive it to be risky.

**Data Source(s):** NSDUH, 2010-12.

**Summary:** During the period 2010-12, Downeast PHD residents ages 18-25 indicated the lowest perception of risk from binge drinking among age groups within the district at 28 percent, while those 26 and older had the highest proportion (45%). Aroostook/Downeast's rates of perception are similar to those of the state. Although not shown, rates of perception of risk from binge drinking among Aroostook/Downeast residents 12 and older have remained relatively unchanged for the past several years.

**Figure 50. Percent of population age 12 or older who perceive a great risk from binge drinking, by Public Health District: 2010-12**



Source: NSDUH


**Indicator Description: PERCEIVED RISK OF REGULAR MARIJUANA USE AMONG YOUTH.** This measure demonstrates the percentage of individuals who perceive a moderate-to-great risk of harm from smoking marijuana regularly.

**Why Indicator is Important:** High school students who do not believe there is moderate to great risk in smoking marijuana regularly are almost eight times as likely to smoke marijuana as their peers who do perceive risk of harm. A similar relationship exists between adult perceptions and consumption.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** Almost six out of ten high school students in Downeast PHD (58%) indicated that there is a moderate-to-great risk of people harming themselves if they smoke marijuana regularly; this was similar to the statewide average of 56 percent.

**Figure 51. Percent of high school students by Public Health District who reported a risk of harm from smoking marijuana regularly: 2009-2013**



Source: MIYHS


**Indicator Description: PERCEIVED RISK OF MARIJUANA USE AMONG MAINERS.** This measure demonstrates the percentage of Mainers over the age of 12 who perceive a risk of harm from smoking marijuana once a month. Because of small sample sizes, survey data from multiple years must be combined in order to produce this estimate.

**Why Indicator is Important:** The perception that using a substance is risky indicates an individual is knowledgeable about health risks and other negative consequences associated with that substance. Perceptions of risk reduce the likelihood that an individual will engage in the behavior.

**Data Source(s):** NSDUH, 2006-08 to 2010-12.

**Summary:** From 2006-08 to 2008-10, the percentage of Aroostook and Downeast PHD<sup>6</sup> residents 12 and older who perceived a great risk from smoking marijuana once a month decreased from 36 percent to 33 percent. Aroostook/Downeast had the highest perception of risk among public health districts in 2008-10.

**Figure 52. Percent of population age 12 or older who perceive a great risk from smoking marijuana once a month by Public Health District: 2006-2008 to 2010-12\***



Source: NSDUH

\*Estimates for Aroostook and Downeast (as single entities) are not available for 2006-08 and 2008-10. Years prior to 2010-12 combine Aroostook and Downeast public health districts.

<sup>6</sup> Due to small sample sizes, Aroostook and Downeast Public Health District (which consists of Washington County and Hancock County) were combined to produce this estimate.


**Indicator Description: PARENTAL ATTITUDES REGARDING MARIJUANA USE.** This indicator reflects how parents felt about their teen using marijuana. Maine parents of teenagers (7<sup>th</sup> thru 12<sup>th</sup> graders) were asked to select the response that best described their attitude about marijuana use by their child. Response options were mutually exclusive.

**Why Indicator is Important:** Parental perceptions and permissive attitudes towards substance use can have a major effect in their child's decision to use. As Maine observes changes in regulations and policies regarding marijuana use; cultural norms and beliefs around use are occurring as well.

**Data Source(s):** Parent Survey, 2013 and 2015

**Summary:** In 2015, 67 percent of parents in Downeast PHD felt it was never okay for their teen to use marijuana; this matched the statewide rate (67%). The rate of disapproval among parents concerning teen marijuana use dropped by 14 percentage points since 2013.

**Figure 53. Parent disapproval regarding their teen using marijuana: 2013 and 2015**






**Indicator Description: PERCEIVED RISK OF PRESCRIPTION DRUG MISUSE AMONG YOUTH.** This measure demonstrates the percentage of individuals who perceive a moderate-to-great risk of harm from taking a prescription drug that was not prescribed to them.

**Why Indicator is Important:** According to the 2013 statewide MIYHS, high school students who do not believe there is moderate-to-great risk misusing prescription drugs are 4.6 times as likely to smoke marijuana as their peers who do perceive risk of harm.

**Data Source(s):** MIYHS, 2009-2013.

**Summary:** In 2011, 85 percent of high school students in Downeast PHD thought misusing prescription drugs was risky; this was on par with the statewide average. That means 15 percent of students did not think misusing prescription drugs was risky.

**Figure 54. Percent of high school students who reported a risk of harm from misusing prescription drugs, by Public Health District: 2009-2013**



Source: MIYHS

## Perceived Enforcement


**Indicator Description: PERCIEVED RISK OF BEING CAUGHT FOR DRINKING ALCOHOL AMONG YOUTH.** This indicator reflects the percentage of high school students who reported that they would be caught by their parents or by police if they drank alcohol.

**Why Indicator is important:** According to the 2013 statewide MIYHS, high school students who believe they would not be caught by their parents are five times as likely to drink in the past month as compared to students who do think they will be caught. Students who believe that they would not be caught by the police are twice as likely to drink alcohol in the past month as those who do think they would be caught.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, 45 percent of Downeast high school students indicated a perceived risk of being caught by their parents for drinking alcohol, which differed little from the state average (43%). Only 16 percent of high school students indicated that they thought they would be caught by the police for drinking alcohol.

**Figure 55. Perceived risk among high school students in Downeast PHD of being caught by parents or police for drinking alcohol: 2009-2011**



Source: MIYHS

**Indicator Description: PERCEIVED RISK OF BEING CAUGHT FOR SMOKING MARIJUANA**


**AMONG YOUTH.** This measure shows the percentage of high school students who reported that they thought they would be caught by police if they smoked marijuana.

**Why Indicator is Important:** According to the statewide 2013 MIYHS, high school students who believe they would be caught by the police are less than half as likely to smoke marijuana as their peers.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** Twenty-four percent of high school students in Downeast PHD indicated that they thought they would be caught by the police if they smoked marijuana, compared to 25 percent statewide who reported this perception. This means that almost three quarters of high school students in Downeast PHD did not think they would be caught by the police if they smoked marijuana.

**Figure 56. Perceived risk among high school students of being caught by police for smoking marijuana, by Public Health District: 2009-2013**



Source: MIYHS

## Mental Health, Suicide and Co-occurring Disorders

---

The relationship between substance use and mental health has been well documented. There are great efforts underway at the Substance Abuse Mental Health Services Administration (SAMHSA) and throughout Maine to better integrate mental health promotion and substance abuse prevention. At the individual level, it is important to know if one exists because the symptoms of each can affect the other; that is, a person who is depressed may abuse alcohol or drugs in an effort to feel better. At the community level, it is important to understand how the prevalence of one interacts with the other so that prevention and intervention efforts can better address the needs of both. The data indicators included below represent the first attempt to collect multiple mental health indicators that can be routinely monitored in relation to substance abuse in hopes that this will lead to better prevention and intervention.

Less than one-fifth of adults in Downeast PHD report having ever been diagnosed with anxiety, a statewide low, and almost one-quarter of high school students felt sad or hopeless every day for two weeks in 2011; more than one in ten high school students reported having considered suicide. The proportion of individuals from Downeast PHD admitted for substance abuse treatment who also have a mental health diagnosis is significantly lower than the statewide rate, but has been gradually increasing since 2009.

## Depression and Anxiety


**Indicator Description: DIAGNOSIS OF ANXIETY AND DEPRESSION AMONG ADULTS.** This indicator examines the percentage of Maine residents age 18 and older who have ever been told by a doctor that they have a depressive or anxiety disorder.

**Why Indicator is Important:** The link between mental health and substance abuse is well documented. Experiencing anxiety or depression is associated with higher rates of substance abuse.

**Data Source(s):** BRFSS, 2011-12 and 2012-13.

**Summary:** The rate of adults in Downeast PHD reporting they have been diagnosed with an anxiety disorder decreased by two percentage points from 2011-12 (18%) to 2012-13 (16%); this was lower than the 2012-13 statewide rate (19%) and lowest among public health districts. Although not shown, in 2012-13, Downeast PHD females were more likely to report having been diagnosed with anxiety when compared to males (20% compared to 10%).


**Figure 57. Percent of adults who have ever been told they have an anxiety disorder, by Public Health District: 2011-12 to 2012-13**



Source: BRFSS

**Summary:** The rate of Downeast PHD adults reporting they had been diagnosed with a depression disorder decreased by two percentage points from 2011 (23%) to 2013 (21%); this was the lowest among the public health districts.

**Figure 58. Percent of adults who have ever been told they have a depression disorder, by Public Health District: 2011-2013**



Source: BRFSS


**Indicator Description: DEPRESSION AMONG YOUTH.** This indicator measures the percentage of high school students reporting they felt sad or hopeless almost every day for two weeks in a row during the past year.

**Why Indicator is Important:** Experiencing depression in the past year is associated with higher rates of substance abuse. According to the 2013 MIYHS, students who reported feeling hopeless or sad for at least two weeks within the past twelve months were almost twice as likely to have used marijuana or to have engaged in binge drinking in the past 30 days, and three times as likely to have misused prescription drugs during the past 30 days. Among youth, depression is also associated with problems with relationships and academic achievement.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, approximately 23 percent (almost one in four) of high school students in Downeast PHD indicated that they felt sad or hopeless every day for two weeks or more in a row during the past year; this was similar to the rate reported by all Maine high school students (23%).

**Figure 59. Felt sad or hopeless almost every day for two weeks or more in a row during the past year, by Public Health District: 2009-2013**



Source: MIYHS

## Suicide and Suicidal Ideation


**Indicator Description: RATE OF SUICIDE DEATHS.** Every death in Maine has a recorded cause. This indicator examines deaths that were classified as a suicide. In this case, a rate per 100,000 of the state population is used to compare the prevalence across certain populations.

**Why Indicator is Important:** Although not the leading cause of death, substance use and abuse is often a factor in suicides. For example, the federal Substance Abuse and Mental Health Services Administration (SAMHSA) has estimated that 23 percent of suicides are attributable to alcohol nationally.

**Data Source(s):** ODRVS, 2013-14

**Summary:** During the two year period 2013-14, Downeast PHD reported a yearly average of 19.6 suicide deaths per 100,000 residents; this was the second highest rate among public health districts. Although not shown, there were 16 suicide deaths in Downeast PHD in 2013 and 18 in 2014.

**Figure 60. Number of suicide deaths per 100,000 residents, by Public Health District: 2013-14**






**Indicator Description: SUICIDAL IDEATION AMONG YOUTH.** This measure examines the percentage of high school students who reported that they seriously considered attempting suicide during the past year.

**Why Indicator is Important:** Suicide is the most tragic consequence of major depressive disorders. Abuse of alcohol or other drugs may increase emotional problems leading to suicidal ideation and suicidal behavior.

**Data Source(s):** MIYHS, 2009-2013; MIYHS 2013 unavailable for Downeast Public Health District due to insufficient sample size.

**Summary:** In 2011, 11 percent of high school students in Downeast PHD considered suicide during the past year, compared to rate reported by high school students statewide (13%).

**Figure 61. Percent of high school students who considered suicide during the past year, by Public Health District: 2009-2013**



Source: MIYHS

## Mental Health and Substance Abuse Co-Occurrence

### Indicator Description: CO-OCCURRING MENTAL HEALTH AND SUBSTANCE ABUSE


**TREATMENT.** This indicator reflects the proportion of treatment admissions for substance abuse where the individual also has a mental health diagnosis.

**Why Indicator is Important:** The link between mental health and substance abuse is well documented. In terms of treatment, it is important to know if one exists because the symptoms of each can affect the other.

**Data Source(s):** WITS, 2010-2014.

**Summary:** In 2014, 48 percent of individuals in Downeast PHD admitted for substance abuse treatment also had a mental health diagnosis; this was considerably lower the statewide rate (58%) and second lowest among public health districts. Downeast PHD has consistently observed some of the lowest rates among public health districts for the past several years.

**Figure 62. Percent of individuals by Public Health District admitted for substance abuse treatment that also had a mental health diagnosis: 2010-2014**



Source: WITS

**Indicator Description: INFORMATION CALLS FOR MENTAL HEALTH AND HUMAN SERVICES.**


2-1-1 Maine is a telephone and internet service that provides information and referrals to health and human services. This indicator reflects the number of calls received by Maine 2-1-1 by the type of service requested.

**Why Indicator is Important:** The data collected from each call provides valuable information serving as a barometer of health and human service needs in the state.

**Data Source(s):** 2-1-1 Maine, 2010-2014

**Summary:** In 2014, Downeast PHD observed 15.8 Maine 2-1-1 referral calls related to mental health services per 10,000 residents; this was much lower than the statewide rate (31.7 calls per 10,000 residents) and second lowest among public health districts. Downeast PHD's rates have remained relatively stable since 2010.

**Figure 63. Number of Maine 2-1-1 referral calls related to mental health services per 10,000 residents, by public health districts: 2010-2014**



Source: 211 Maine, 2010-2014

## Treatment Admissions for Substance Abuse

---

Substance abuse treatment admissions are an indicator of how many people *receive treatment* for a substance abuse problem. These admissions can be voluntary, but they can also be court-ordered. Treatment admission data should not be used as an indicator of the magnitude of the problems related to substance abuse. Rather, treatment should be seen as a major consequence stemming from substance use and one that requires many resources.

Mainers continue to seek out treatment for abuse involving a wide array of substances besides alcohol; in 2014 3,589 clients were admitted for alcohol as the primary substance. This was followed by synthetic opioids (2,663) and heroin (2,538).

In 2014, treatment admissions for synthetic opioids dropped slightly after a dramatic increase in 2011, but remained significantly higher than alcohol-related treatment admissions; the rate was highest in the state for such admissions. Although proportions of primary admissions related to heroin/morphine are lower than the state average, Downeast PHD observed a notable increase from 2012 to 2014. Downeast PHD has a slightly higher proportion of secondary admissions related to marijuana as compared to the statewide average, while synthetic opioids are a close second in this respect.

## Treatment Admissions


**Indicator Description: PRIMARY TREATMENT ADMISSIONS.** This measure reflects substance abuse treatment admissions by licensed substance abuse treatment agencies/providers. A “primary” substance is identified during the treatment admissions process based on use patterns (e.g., frequency, duration, quantity) and the risk(s) posed to the individual. The analysis excludes admissions for shelter/detoxification services.

**Why Indicator is Important:** The number of substance abuse treatment admissions is bound by both the need and the capacity for treatment. Treatment admissions data are not a good indicator of substance use, abuse or dependence, but provide an indication of service usage and the impact of substance use on the behavioral healthcare system.

**Data Source(s):** WITS, 2010-2014.

**Summary:** In 2014, 35 percent of all primary treatment admissions in Downeast PHD were related to synthetic opioids<sup>7</sup> (35%), followed by alcohol (33%) and marijuana (11%). The majority of primary admissions for treatment in Downeast PHD have been related to synthetic opioids since 2010. Primary admissions related to synthetic opioids in Downeast PHD have been steadily decreasing since 2011.

**Figure 64. Primary drug admissions for in Downeast PHD, by drug type: 2010-2014**




Source: WITS

<sup>7</sup> “Synthetic opioids” excludes methadone and buprenorphine.

**Summary:** In 2014, the proportion of primary treatment admissions for synthetic opioids in Downeast PHD was 10 percentage points higher than the statewide average (35% compared to 25%), while the rate for heroin was 7 percentage points lower than the state (17% compared to 24%).


**Figure 65. Primary treatment admissions in Downeast PHD: 2014**



Source: WITS

**Summary:** In 2014, Downeast PHD was below the average (31%) among public health districts of primary treatment admissions due to alcohol (34%).


**Figure 66. Primary admissions related to alcohol, by Public Health District and drug type: 2010-2014**



Source: WITS

**Summary:** In 2014, Downeast PHD had the highest rate of primary treatment admissions related to synthetic opioids. The proportion of primary admissions related to synthetic opioids decreased by 13 percentage points since 2011 (from 48% to 35%).

**Figure 67. Primary admissions related to synthetic opioids, by Public Health District: 2010-2014**




Source: WITS



**Summary:** Downeast PHD has observed a steady increase in the proportion of primary admissions due to heroin, since 2010 admissions have increased 13 percentage points.


**Figure 68. Primary admissions related to heroin/morphine, by Public Health District: 2010-2014**



Source: WITS

**Summary:** From 2012 to 2014, Downeast PHD observed an average proportion of primary treatment admissions related to marijuana. Downeast PHD's rate remained close to the statewide average from 2010-14.

**Figure 69. Primary admissions related to marijuana, by Public Health District and drug type: 2010-2014**



Source: WITS


**Indicator Description: SECONDARY TREATMENT ADMISSIONS.** This measure reflects substance abuse treatment admissions by licensed substance abuse treatment agencies/providers. A “secondary” substance is identified during the admissions process as one used by the individual and for which treatment may be received, but it is not the primary substance for which treatment was sought. The analysis excludes admissions for shelter/detoxification services.

**Why Indicator is Important:** The number of substance abuse treatment admissions is bound by both the need and the capacity for treatment. Treatment admission data are not a good indicator of substance use, abuse or dependence, but provide an indication of service usage and the impact of substance use on the behavioral healthcare system.

**Data Source(s):** WITS, 2010-2014.

**Summary:** In 2014, 29 percent of secondary treatment admissions in Downeast PHD were for marijuana, followed by synthetic opioids (28%) and then Heroin (14%). While secondary admissions related to marijuana, alcohol and synthetic opioids have remained relatively stable over the past several years, Heroin has begun an increasing trend.


**Figure 70. Secondary treatment admissions in Downeast PHD: 2010-2014**



Source: WITS

**Summary:** In 2014, Downeast PHD had a greater proportion of secondary admissions related to alcohol (12%), heroin/morphine (14%), and synthetic opioids (28%) compared to secondary treatment admissions statewide. While Downeast PHD's proportions of secondary treatment admissions related to marijuana and methadone were on par with those of the state, it observed much lower rates regarding benzodiazepines and cocaine/crack.

**Figure 71. Secondary treatment admissions in Downeast PHD, by drug type:**  
**2014**



Source: WITS



## Appendix: Data Sources

---

This report includes data that was gathered from a number of data sources. A detailed description of each source is provided below, consisting of information about the data included in each source, and retrieval or contact information. The report includes data that were available through May 2015.

There are multiple purposes for this report. One is to provide a snapshot of the most recent data regarding substance abuse, while another is to examine trends over time. Therefore, each indicator may have multiple sources of data that are included. While each indicator provides a unique and important perspective on drug use in Maine, none should individually be interpreted as providing a full picture of drug trends in Maine. In particular, the percentages and figures from one data source do not always align with the data and percentages from a similar source. Older data are often included in order to examine an indicator among a specific population or to find trends over time. When discussing rates of prevalence, however, the user should rely upon the most recent data source available.

### Description of Data Sources

**Behavioral Risk Factor Surveillance System (BRFSS).** The BRFSS is a national survey administered on an ongoing basis by the National Centers for Disease Control and Prevention (CDC) to adults in all 50 states and several districts and territories. The instrument collects data on adult risk behaviors, including alcohol abuse. BRFSS defines heavy drinking as adult men having more than two drinks per day and adult women having more than one drink per day, and binge drinking as males having five or more drinks on one occasion and females having four or more drinks on one occasion. The most recent data available are from 2012. **Due to methodological changes in weighting and sampling, data prior to 2011 cannot be trended with more current data.** In addition, in some cases years were combined to yield more stable estimates. Contact: Melissa Damren, Maine BRFSS Coordinator; [melissa.damren@maine.gov](mailto:melissa.damren@maine.gov); (207) 287-1420.

**Maine Department of Public Safety (DPS), Uniform Crime Reports (UCR).** UCR data include drug and alcohol arrests. Drug arrests include sale and manufacturing as well as possession of illegal substances. Liquor arrests include all liquor law violations. OUI arrests are arrests for operating a motor vehicle under the influence of a controlled substance. DPS data are now available from 2013. Arrest data may reflect differences in resources or focus of law enforcement efforts, so may not be directly comparable from year to year. Retrieval: [http://www.maine.gov/dps/cim/crime\\_in\\_maine/cim.htm](http://www.maine.gov/dps/cim/crime_in_maine/cim.htm)

**Maine Department of Public Safety (DPS), Liquor Licensing and Compliance.** DPS issues and renews licenses for the manufacture, importation, storage, transportation and sale of all liquor and administers those laws relating to licensing and the collection of taxes on malt liquor and wine. DPS maintains a list of all active licenses that can be accessed online.

Retrieval: [http://www.maine.gov/dps/liqr/active\\_licenses.htm](http://www.maine.gov/dps/liqr/active_licenses.htm)

**Maine Department of Public Safety (DPS), Bureau of Highway Safety (BHS), Maine Department of Transportation (MDOT).** The Bureau of Highway Safety is responsible for tracking all fatalities that occur on Maine's highways and reporting this information through the Fatal Analysis Reporting System (FARS). The data represented provides information on highway crashes and fatalities. Much of this information is gathered from our FARS system, which records data on fatal crashes in Maine for input into a larger national record-keeping system of statistical data. FARS data is also used by BHS and the Maine State Police to analyze enforcement priorities and schedules. Impaired driving is one of the most serious traffic risks facing the nation, killing thousands every year. Contact: Duane Brunell, Safety Performance Analysis Manager; [duane.brunell@maine.gov](mailto:duane.brunell@maine.gov); (207) 624-3278.

**Maine Drug Enforcement Agency (MDEA).** The MDEA, through its eight regional multi-jurisdictional task forces, is the lead state agency in confronting drug trafficking crime. This indicator differs from the Uniform Crime Report drug-related arrest data in that it only tracks MDEA efforts and does not encompass all activity within Maine law enforcement agencies. The data included in this report represents those arrested for a drug offense but does not indicate what other drug(s) may have been seized. For example, a person may be arrested for the sale of cocaine but also be in possession of oxycodone and marijuana. It is important to note that arrests and multi-jurisdictional drug enforcement are resource-dependent; such funds fluctuate from year to year, and must be reallocated to combat highest priority threats. Contact: Roy E. McKinney, Director; [roy.e.mckinney@maine.gov](mailto:roy.e.mckinney@maine.gov); (207) 626-3852.

**Maine Emergency Medical Services (EMS).** Maine EMS is a bureau within the Maine Department of Public Safety (DPS) and is responsible for the coordination and integration of all state activities concerning Emergency Medical Services and the overall planning, evaluation, coordination, facilitation and regulation of EMS systems. EMS collects data statewide from the 272 licensed ambulance and non-transporting services. It is mandated that services submit an electronic patient care report to Maine EMS within one business day of patient contact. Data are compiled upon request. Contact: Jon Powers, Maine Emergency Medical Services; [jon.powers@maine.gov](mailto:jon.powers@maine.gov); (207) 626-3860.

**Maine Integrated Youth Health Survey (MIYHS).** The MIYHS is a statewide survey administered biennially through a collaborative partnership by the Maine Office of Substance Abuse (OSA) the Maine Center for Disease Control and Prevention and the Maine department of Education to students in grades 5 through 12. The survey collects information on student substance use, risk factors related to substance use, as well as consequences, perceptions and social risk factors related to substances, and collects information on many other health factors. As of the date of this report, the most recent data available are from 2013. Due to changes in the survey administration and structure, the new survey data cannot be trended with the Maine Youth Drug and Alcohol Survey (MYDAUS). In addition, **due to a low response rate, the Downeast public health district does not have estimates for 2013.** Contact: Stephen Corral, Substance

Abuse Program Specialist, Office of Substance Abuse, [stephen.corral@maine.gov](mailto:stephen.corral@maine.gov); (207) 287-2964.

**Maine Health Data Organization (MHDO).** MHDO data includes all inpatient admissions to all hospitals in Maine for calendar years 2010 and 2011. Data categories created by the authors include alcohol, opioids, illegal drugs, and pharmaceuticals. All drug categories include intoxication, abuse, dependence, and poisoning cases related to the drug. The opiate category includes methadone, heroin, and opiates. The illegal drug category includes crack/cocaine, cannabis, and hallucinogens. The pharmaceuticals category includes all other non-opioid medications (including stimulants and depressants). Contact: Maine Health Data Organization (MHDO), [lisa.parker@maine.gov](mailto:lisa.parker@maine.gov); (207) 287-3225.

**Marci Sorg, Margaret Chase Smith Policy Center at University of Maine, Office of the Chief Medical Examiner.** The Maine Office of the Chief Medical Examiner maintains records of all deaths associated with drug overdose. Drug categories include methadone, cocaine, benzodiazepines, oxycodone and heroin/morphine. The death data are compiled on an annual basis and must be finalized prior to release and so are not available to track changes that may occur over shorter time frames. Contact: Dr. Marcella Sorg, Director, Rural Drug & Alcohol Research Program, Margaret Chase Smith Policy Center, University of Maine; [marcella\\_sorg@umit.maine.edu](mailto:marcella_sorg@umit.maine.edu); (207) 581-2596.

**National Survey on Substance Use and Health (NSDUH).** The NSDUH is a national survey administered annually by the Substance Abuse and Mental Health Services Administration (SAMHSA) to youth grades 6 through 12 and adults ages 18 and up. The instrument collects information on substance use and health at the national, regional and state levels. The advantage of NSDUH is that it allows comparisons to be made across the lifespan (that is, ages 12 and up). However, NSDUH is not as current as other data sources; as of this report, data at the substate level are available from 2008-2010. Older data are included for trending and comparative purposes. NSDUH defines Illicit Drugs as marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used non-medically; Binge Alcohol Use as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least one day in the past 30 days; Dependence or abuse based on definitions found in the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV); and Serious Mental Illness (SMI) as a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. Retrieval: <https://nsduhweb.rti.org/>

**Northern New England Poison Center (NNEPC).** The Northern New England Poison Center provides services to Maine, New Hampshire, and Vermont. A poisoning case represents a single individual's contact with a potentially toxic substance. Intentional poisoning includes those related to substance abuse, suicide and misuse. Data include the number of confirmed cases where exposures are judged to be substance abuse-related (i.e., an individual's attempt to get



high). NNEPC collects detailed data on specific substances involved in poisonings, including the categories of stimulants/street drugs, alcohol, opioids, asthma/cold and cough, benzodiazepines, antidepressants, and pharmaceuticals, as well as other substances. The category of stimulants/street drugs includes marijuana and other cannabis, amphetamine and amphetamine-like substances, cocaine (salt and crack), amphetamine/dextroamphetamine, caffeine tablets/capsules, ecstasy, methamphetamine, GHB, and other/unknown stimulants/street drugs. The category alcohol includes alcohol-containing products such as mouthwash. The opioid category includes Oxycodone, Hydrocodone, buprenorphine, methadone, tramadol, morphine, propoxyphene, codeine, hydromorphone, stomach opioids, Meperidine (Demerol), heroin, Fentanyl, and other/unknown opioids. The asthma/cold and cough category includes eye, ear, nose, and throat medications. Data available from the poison center are reported on a continual daily basis and are included through December 2013. These data are only reflective of cases in which the Poison Center was contacted. Contact: Karen Simone, Director, Northern New England Poison Center; [simonk@mmc.org](mailto:simonk@mmc.org); (207) 662-7221.

**Office of Child and Family Services (OCFS), Maine Automated Child Welfare Information System (MACWIS).** The Office of Child and Family Services (OCFS) supports Maine's children and their families by providing Children's Behavioral Health, Child Welfare, Early Childhood, and Community services. The Maine Child Welfare Information System (MACWIS) serves as the single repository for all Maine child welfare information to assist Office of Child and Family Services (OCFS) workers in the recording, tracking, and processing of child welfare functions. MACWIS is the single repository for all electronic child welfare information. It actively manages 850,000 identified persons and 28,525 resources in the system; [lori.geiger@maine.gov](mailto:lori.geiger@maine.gov); (207)-624-7911.

**Prescription Monitoring Program (PMP).** PMP maintains a database of all transactions for class C-II through C-IV drugs dispensed in the state of Maine. Drug categories used in this report include narcotics, tranquilizers, stimulants, and other prescriptions. Other prescriptions include those drugs that are not classified as narcotics, tranquilizers or stimulants, including products such as endocrine and metabolic drugs, analgesics and anesthetics, gastrointestinal agents, and nutritional products. The counts included in this report represent the quantity dispensed through prescriptions filled between 2009 and 2013. Contact: John Lipovsky, PMP Coordinator, Substance Abuse and Mental Health Services; [john.lipovsky@maine.gov](mailto:john.lipovsky@maine.gov); (207) 287-3363.

**WITS Substance Abuse Treatment Data System (WITS).** WITS is a statewide database that includes information about clients admitted to treatment in SAMHS-funded facilities. Analyses in this report are based on clients' reported primary, secondary and tertiary drug(s) of choice as well as other demographic and background information that is collected at intake. Drug categories included in this report are alcohol, marijuana, cocaine, heroin, synthetic opiates, methadone/buprenorphine and benzodiazepines. Contact: Johanna Buzzell, Substance Abuse and Mental Health Services; [johanna.buzzell@maine.gov](mailto:johanna.buzzell@maine.gov); (207) 287-6337.

**U.S. Census Bureau.** The U.S. Census provides summary profiles showing frequently requested data items from various Census Bureau programs. Profiles are available for all states and

counties, and for cities and towns with more than 25,000 people. Data are updated no less than annually. Retrieval for Maine census data: <http://quickfacts.census.gov/qfd/states/23000.html>